

Your source for the latest research news

# **Science News**

from research organizations

# How COVID-19 spread has been contained by travel bans

New computer modelling could play a big part in exit strategies and lifting air travel restrictions

Date: May 5, 2020

Source: Taylor & Francis Group

Summary: Millions more people across the EU could have contracted COVID-19 had strict international travel

bans not been implemented, shows a new report by computer modelling experts.

Share: **f y p** in **x** 

#### **FULL STORY**

Millions more people across the EU could have contracted COVID-19 had strict international travel bans not been implemented, shows a new report by computer modelling experts at Stanford University.

Using a newly developed mathematical epidemiology simulation, the study, published in *Computer Methods in Biomechanics and Biomedical Engineering*, predicts the huge impact that limiting air travel across the 27 EU nations had on restricting the spread of the disease.

The simulation can show live estimated figures for the growth of spread for each country if we were to remove travel bans today. The images above show how 0.2% of some populations could have become infected by 20th April (when the study was written, 5 April), however these figures change daily.

This new model could now play a vital part in establishing politicians' exit strategies, with the team able to virtually lift travel restrictions between individual communities, states, or countries, to explore the potential gradual changes in spreading patterns and outbreak dynamics.

"There is a well-reasoned fear that easing of current (travel restriction) measures, even slightly, could trigger a new outbreak and accelerate the spread to an unmanageable degree," lead author Ellen Kuhl, Professor of Mechanical Engineering at Standford University comments.

"Global network mobility models, combined with local epidemiology models, can provide valuable insight into different exit strategies. Our results demonstrate that mathematical modelling can provide guidelines for political decision making with the ultimate goal to gradually return to normal while keeping the rate of new COVID-19 infections steady and manageable," says Kevin Linka, lead author and postdoctoral researcher in Dr. Kuhl's group.

From its European origin in Italy, the novel coronavirus spread rapidly via the strongest network connections to Germany, Spain, and France, while slowly reaching the less connected countries, Estonia, Slovakia, and Slovenia.

Currently the levels of population known to be infected with the disease varies from country to country, however as of April 18, with flight being reduced by 89% in Germany, 93% in France, 94% in Italy, and 95% in Spain (Eurostat 2020), the graphs in this study show how the spread has been contained.

"Strikingly, our results suggest that the emerging pattern of the COVID-19 outbreak closely followed global mobility patterns of air passenger travel," confirms Professor Kuhl, whose model can also predict the emerging global diffusion pattern of a pandemic at the early stages of the outbreak.

"Our results suggest that an unconstrained mobility would have significantly accelerated the spreading of COVID-19, especially in Central Europe, Spain, and France."

Unfortunately, the model also confirms how travel bans were introduced too late to stop the Europe-wide outbreak altogether.

"A recent study based on a global metapopulation disease transmission model for the COVID-19 outbreak in China has shown that the Wuhan travel ban essentially came too late, at a point where most Chinese cities had already received many infected travellers (Chinazzi et al. 2020). Our study shows a similar trend for Europe, where travel restrictions were only implemented a week after every country had reported cases of COVID-19 (European Centre for Disease Prevention and Control 2020).

"As a natural consequence, unfortunately, no European country was protected from the outbreak," Professor Kuhl, who is the Robert Bosch Chair of Mechanical Engineering at Standford added.

The first official case of COVID-19 in Europe was reported in France on January 24, 2020, followed by Germany and Finland only three and five days later. Within only six weeks, all 27 countries of the European Union were affected, with the last cases reported in Malta, Bulgaria, and Cyprus on March 9, 2020. At this point, there were 13,944 active cases within the European Union and the number of active cases doubled every three to four days (European Centre for Disease Prevention and Control 2020).

Dr Kuhl adds that although air travel is certainly not the only determinant of the outbreak dynamics, their findings indicate that "mobility is a strong contributor to the global spreading of COVID-19." This is becoming especially important now that many countries are beginning to lift their travel restrictions in an attempt to gradually return to normal.

Other limitations highlighted -- like any infectious disease model -- include the simulation being subject to data uncertainties from differences in testing, inconsistent diagnostics, incomplete counting, and delayed reporting across all countries.

### **Story Source:**

Materials provided by Taylor & Francis Group. Note: Content may be edited for style and length.

### Journal Reference:

 Kevin Linka, Mathias Peirlinck, Francisco Sahli Costabal, Ellen Kuhl. Outbreak dynamics of COVID-19 in Europe and the effect of travel restrictions. Computer Methods in Biomechanics and Biomedical Engineering, 2020; 1 DOI: 10.1080/10255842.2020.1759560

### Cite This Page:

MLA

**APA** 

Chicago

Taylor & Francis Group. "How COVID-19 spread has been contained by travel bans: New computer modelling could play a big part in exit strategies and lifting air travel restrictions." ScienceDaily. ScienceDaily, 5 May 2020. <a href="https://www.sciencedaily.com/releases/2020/05/200504191018.htm">www.sciencedaily.com/releases/2020/05/200504191018.htm</a>.

#### **RELATED STORIES**

US Modelling Study Estimates Impact of School Closures for COVID-19 on US Health-Care Workforce and Associated Mortality

Apr. 4, 2020 — US policymakers considering physical distancing measures to slow the spread of COVID-19 face a difficult trade-off between closing schools to reduce transmission and new cases, and potential ... **read more** »

Effectiveness of Travel Bans -- Readily Used During Infectious Disease Outbreaks -- Mostly Unknown Feb. 13, 2020 — While travel bans are frequently used to stop the spread of an emerging infectious disease, a new study of published research found that the effectiveness of travel bans is mostly ... **read more** »

Feel Anxious? Have Trouble sleeping? You May Be Traveling for Business Too Often

Jan. 8, 2018 — People who travel for business two weeks or more a month report more symptoms of anxiety and depression and are more likely to smoke, be sedentary and report trouble sleeping than those who travel ... **read more** »

Infectious Disease Spread Is Fueled by International Trade

Dec. 22, 2015 — International trade and travel has literally opened up new vistas for humans, ranging from travel to exotic places to enjoying the products and services of those distant lands. But along with ... **read more** »

#### FROM AROUND THE WEB

Below are relevant articles that may interest you. ScienceDaily shares links with scholarly publications in the TrendMD network and earns revenue from third-party advertisers, where indicated.

SARS-CoV-2 Experience Will Help Agency Strike Right Balance With LDTs, FDA Says 🗹

Kelsy Ketchum, 360Dx, 2020

Coronavirus in the Crescent City <a></a>

Ciara Curtin, GenomeWeb, 2020

GeneFirst, Rapid Diagnostics Validating Coronavirus Assay for Point-of-Care Platform 🗹

Tony Fong, GenomeWeb, 2020

Merck, Promega Collaborate to Develop MSI CDx for Keytruda 🗹

Precision Oncology News, 2019

Coronavirus Dx Emergency Use Authorizations Progressing Rapidly Despite Criticism 🗹

Madeleine Johnson, 360Dx, 2020

Transiting (in) Shanghai: High-skilled professionals from Spain to China 🗹

Masdeu Torruella et al., Transitions: Journal of Transient Migration

Oncologists See Clinical Trial Disruptions as Coronavirus Pandemic Continues 🗹

Precision Oncology News, 2020

German Research Project Shows Feasibility of SARS-CoV-2 Testing Using Hologic LDT Functionality <a>Idea</a> O'Connor, 360Dx, 2020





# **Free Subscriptions**

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

- RSS Feeds

## **Follow Us**

Keep up to date with the latest news from ScienceDaily via social networks:

- **f** Facebook
- in LinkedIn

### Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

- Leave Feedback
- Contact Us

About This Site | Staff | Reviews | Contribute | Advertise | Privacy Policy | Editorial Policy | Terms of Use

Copyright 2020 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners. Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.

Do Not Sell My Personal Information