

## Science News

*from research organizations*

### Spitzer telescope reveals the precise timing of a black hole dance


*Date:* April 28, 2020

*Source:* NASA/Jet Propulsion Laboratory

*Summary:* Scientists have finally figured out the precise timing of a complicated dance between two enormous black holes, revealing hidden details about the physical characteristics of these mysterious cosmic objects.

*Share:* [f](#) [t](#) [p](#) [in](#) [✉](#)

#### FULL STORY



This image shows two massive black holes in the OJ 287 galaxy. The smaller black hole orbits the larger one, which is also surrounded by a disk of gas. When the smaller black hole crashes through the disk, it produces a flare brighter than 1 trillion stars.

*Credit: NASA/JPL-Caltech*

Black holes aren't stationary in space; in fact, they can be quite active in their movements. But because they are completely dark and can't be observed directly, they're not easy to study. Scientists have finally figured out the precise timing of a complicated dance between two enormous black holes, revealing hidden details about the physical characteristics of these mysterious cosmic objects.

The OJ 287 galaxy hosts one of the largest black holes ever found, with over 18 billion times the mass of our Sun. Orbiting this behemoth is another black hole with about 150 million times the Sun's mass. Twice every 12 years, the smaller black hole crashes through the enormous disk of gas surrounding its larger companion, creating a flash of light brighter than a trillion stars -- brighter, even, than the entire Milky **Way** galaxy. The light takes 3.5 billion years to reach Earth.

But the smaller black hole's orbit is oblong, not circular, and it's irregular: It shifts position with each loop around the bigger black hole and is tilted relative to the disk of gas. When the smaller black hole crashes through the disk, it creates two expanding bubbles of hot gas that move away from the disk in opposite directions, and in less than 48 hours the system appears to quadruple in brightness.

Because of the irregular orbit, the black hole collides with the disk at different times during each 12-year orbit. Sometimes the flares appear as little as one year apart; other times, as much as 10 years apart. Attempts to model the orbit and predict when the flares would occur took decades, but in 2010, scientists created a model that could predict their occurrence to within about one to three weeks. They demonstrated that their model was correct by predicting the appearance of a flare in December 2015 to within three weeks.

Then, in 2018, a group of scientists led by Lankeswar Dey, a graduate student at the Tata Institute of Fundamental Research in Mumbai, India, published a paper with an even more detailed model they claimed would be able to predict the timing of future flares to within four hours. In a new study published in the *Astrophysical Journal Letters*, those scientists report that their accurate prediction of a flare that occurred on July 31, 2019, confirms the model is correct.

The observation of that flare almost didn't happen. Because OJ 287 was on the opposite side of the Sun from Earth, out of view of all telescopes on the ground and in Earth orbit, the black hole wouldn't come back into view of those telescopes until early September, long after the flare had faded. But the system was within view of NASA's Spitzer Space Telescope, which the agency retired in January 2020.

After 16 years of operations, the spacecraft's orbit had placed it 158 million miles (254 million kilometers) from Earth, or more than 600 times the distance between Earth and the Moon. From this vantage point, Spitzer could observe the system from July 31 (the same day the flare was expected to appear) to early September, when OJ 287 would become observable to telescopes on Earth.

"When I first checked the visibility of OJ 287, I was shocked to find that it became visible to Spitzer right on the day when the next flare was predicted to occur," said Seppo Laine, an associate staff scientist at Caltech/IPAC in Pasadena, California, who oversaw Spitzer's observations of the system. "It was extremely fortunate that we would be able to capture the peak of this flare with Spitzer, because no other human-made instruments were capable of achieving this feat at that specific point in time."

## **Ripples in Space**

Scientists regularly model the orbits of small objects in our solar system, like a comet looping around the Sun, taking into account the factors that will most significantly influence their motion. For that comet, the Sun's gravity is usually the dominant force, but the gravitational pull of nearby planets can change its path, too.

Determining the motion of two enormous black holes is much more complex. Scientists must account for factors that might not noticeably impact smaller objects; chief among them are something called gravitational waves. Einstein's theory of general relativity describes gravity as the warping of space by an object's mass. When an object moves through space, the distortions turn into waves. Einstein predicted the existence of gravitational waves in 1916, but they weren't observed directly until 2015 by the Laser Interferometer Gravitational Wave Observatory (LIGO).

The larger an object's mass, the larger and more energetic the gravitational waves it creates. In the OJ 287 system, scientists expect the gravitational waves to be so large that they can carry enough energy away from the system to measurably alter the smaller black hole's orbit -- and therefore timing of the flares.

While previous studies of OJ 287 have accounted for gravitational waves, the 2018 model is the most detailed yet. By incorporating information gathered from LIGO's detections of gravitational waves, it refines the window in which a flare is expected to occur to just 1 1/2 days.

To further refine the prediction of the flares to just four hours, the scientists folded in details about the larger black hole's physical characteristics. Specifically, the new model incorporates something called the "no-hair" theorem of black holes.

Published in the 1960s by a group of physicists that included Stephen Hawking, the theorem makes a prediction about the nature of black hole "surfaces." While black holes don't have true surfaces, scientists know there is a boundary around them beyond which nothing -- not even light -- can escape. Some ideas posit that the outer edge, called the event horizon, could be bumpy or irregular, but the no-hair theorem posits that the "surface" has no such features, not even hair (the theorem's name was a joke).

In other words, if one were to cut the black hole down the middle along its rotational axis, the surface would be symmetric. (The Earth's rotational axis is almost perfectly aligned with its North and South Poles. If you cut the planet in half along that axis and compared the two halves, you would find that our planet is mostly symmetric, though features like oceans and mountains create some small variations between the halves.)

### **Finding Symmetry**

In the 1970s, Caltech professor emeritus Kip Thorne described how this scenario -- a satellite orbiting a massive black hole -- could potentially reveal whether the black hole's surface was smooth or bumpy. By correctly anticipating the smaller black hole's orbit with such precision, the new model supports the no-hair theorem, meaning our basic understanding of these incredibly strange cosmic objects is correct. The OJ 287 system, in other words, supports the idea that black hole surfaces are symmetric along their rotational axes.

So how does the smoothness of the massive black hole's surface impact the timing of the smaller black hole's orbit? That orbit is determined mostly by the mass of the larger black hole. If it grew more massive or shed some of its heft, that would change the size of smaller black hole's orbit. But the distribution of mass matters as well. A massive bulge on one side of the larger black hole would distort the space around it differently than if the black hole were symmetric. That would then alter the smaller black hole's path as it orbits its companion and measurably change the timing of the black hole's collision with the disk on that particular orbit.

"It is important to black hole scientists that we prove or disprove the no-hair theorem. Without it, we cannot trust that black holes as envisaged by Hawking and others exist at all," said Mauri Valtonen, an astrophysicist at University of Turku in Finland and a coauthor on the paper.

Spitzer science data continues to be analyzed by the science community via the Spitzer data archive located at the Infrared Science Archive housed at IPAC at Caltech in Pasadena. JPL managed Spitzer mission operations for NASA's Science Mission Directorate in Washington. Science operations were conducted at the Spitzer Science Center at IPAC at Caltech. Spacecraft operations were based at Lockheed Martin Space in Littleton, Colorado. Caltech manages JPL for NASA.

For more information about Spitzer, visit:

- <https://www.nasa.gov/spitzer>
- <http://www.spitzer.caltech.edu/>

---

### Story Source:

Materials provided by **NASA/Jet Propulsion Laboratory**. *Note: Content may be edited for style and length.*

---

### Related Multimedia:

- YouTube video: [Timing of Black Hole Dance Revealed by NASA Spitzer Space Telescope](#)
- 

### Journal Reference:

1. Seppo Laine, Lankeswar Dey, Mauri Valtonen, A. Gopakumar, Stanislaw Zola, S. Komossa, Mark Kidger, Pauli Pihajoki, José L. Gómez, Daniel Caton, Stefano Ciprini, Marek Drozd, Kosmas Gazeas, Vira Godunova, Shirin Haque, Felix Hildebrandt, Rene Hudec, Helen Jermak, Albert K. H. Kong, Harry Lehto, Alexios Liakos, Katsura Matsumoto, Markus Mugrauer, Tapio Pursimo, Daniel E. Reichart, Andrii Simon, Michal Siwak, Eda Sonbas. **Spitzer Observations of the Predicted Eddington Flare from Blazar OJ 287**. *The Astrophysical Journal*, 2020; 894 (1): L1 DOI: 10.3847/2041-8213/ab79a4
- 

### Cite This Page:

MLA

APA

Chicago

---

NASA/Jet Propulsion Laboratory. "Spitzer telescope reveals the precise timing of a black hole dance." ScienceDaily. ScienceDaily, 28 April 2020. <[www.sciencedaily.com/releases/2020/04/200428223728.htm](http://www.sciencedaily.com/releases/2020/04/200428223728.htm)>.

### RELATED STORIES

---

#### Hiding Black Hole Found

Feb. 28, 2019 — Astronomers have detected a stealthy black hole from its effects on an interstellar gas cloud. This intermediate mass black hole is one of over 100 million quiet black holes expected to be lurking in ...

#### Listening for Gravitational Waves Using Pulsars

Nov. 15, 2017 — When galaxies collide, their central black holes tend to spiral toward each other, releasing gravitational waves in their cosmic dance. To explore this uncharted area of gravitational wave science, ...

#### Tail of Stray Black Hole Hiding in the Milky Way

Feb. 2, 2017 — By analyzing the gas motion of an extraordinarily fast-moving cosmic cloud in a corner of the Milky Way, astronomers found hints of a wandering black hole hidden in the cloud. This result marks the ...

#### Funky Light Signal from Colliding Black Holes Explained

Sep. 18, 2015 — Entangled by gravity and destined to merge, two candidate black holes in a distant galaxy appear to be locked in an intricate dance. Researchers using data from NASA's Galaxy Evolution Explorer ...

*Below are relevant articles that may interest you. ScienceDaily shares links with scholarly publications in the TrendMD network and earns revenue from third-party advertisers, where indicated.*

### **Cosmological Implications of the Planck Aether Model for a Unified Field Theory**

Zeitschrift für Naturforschung A, 2014

### **Emission of Gravitational Waves from a Magnetohydrodynamic Dynamo in the Center of the Sun**

Zeitschrift für Naturforschung A, 2015

### **Statistical Mechanical Interpretation of Black Hole Entropy**

Zeitschrift für Naturforschung A, 2014

### **Gravity beyond Einstein? Part I: Physics and the Trouble with Experiments**

Jochem Hauser et al., Zeitschrift für Naturforschung A, 2017

### **An Intuitive Understanding of Black-Hole Evaporation by Viewing it in Terms of More Familiar Quantum-Field Effects in Flat Spacetime**

Zeitschrift für Naturforschung A, 2014

### **The Exchange-Correlation Field Effect over the Magnetoacoustic-Gravitational Instability in Plasmas**

A. Rasheed et al., Zeitschrift für Naturforschung A, 2017

Ettl et al., Breast Cancer Research, 2020

### **The Electronic Structure of Closed Shell Metallocenes in the Ground State and in the Cationic Hole-States**

Zeitschrift für Naturforschung A, 2014

---

Powered by **TREND MD**

## **Free Subscriptions**

---

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

 Email Newsletters

 RSS Feeds

## **Follow Us**

---

Keep up to date with the latest news from ScienceDaily via social networks:

 Facebook

 Twitter

## Have Feedback?

---

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

 [Leave Feedback](#)

 [Contact Us](#)

[About This Site](#) | [Staff](#) | [Reviews](#) | [Contribute](#) | [Advertise](#) | [Privacy Policy](#) | [Editorial Policy](#) | [Terms of Use](#)

Copyright 2020 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners.

Content on this website is for information only. It is not intended to provide medical or other professional advice.

Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners.

Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.